

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

Name:	D.O.B.: PLACE PICTURE	
Allergy to:	HERE	
Weight:Ibs. Asthma: [] Yes (higher risk fo	r a severe reaction) [] No	
NOTE: Do not depend on antihistamines or inhalers ((bronchodilators) to treat a severe reaction. USE EPINEPHRINE.	

Extremely reactive to the following foods: THEREFORE: 1 If checked, give epinephrine immediately for ANY symptoms if the allergen was likely eaten. [] If checked, give epinephrine immediately if the allergen was definitely eaten, even if no symptoms are noted.

FOR ANY OF THE FOLLOWING:

SEVERE SYMPTOMS



Short of breath, wheezing, repetitive cough



HEART

Pale, blue, faint, weak pulse, dizzy



THROAT

Tight, hoarse, trouble breathing/ swallowing



MOUTH

Significant swelling of the tongue and/or lips

OR A

COMBINATION of symptoms

from different

body areas.



Many hives over body, widespread redness



Repetitive vomiting, severe diarrhea



Feeling anxiety, confusion



OTHER



something bad is about to happen,









INJECT EPINEPHRINE IMMEDIATELY.

- 2. Call 911. Tel! them the child is having anaphylaxis and may need epinephrine when they arrive.
- Consider giving additional medications following epinephrine:
 - Antihistamine
 - Inhaler (bronchodilator) if wheezing
- Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport them to ER even if symptoms resolve. Person should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS









NOSE

Itchy/runny nose, sneezing

MOUTH Itchy mouth

A few hives. Mild nausea/

mild itch

discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

- 1. Antihistamines may be given, if ordered by a healthcare provider.
- 2. Stay with the person; alert emergency contacts.
- 3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES


oinephrine Brand:		
oinephrine Dose: []	0.15 mg IM	[] 0.3 mg IM
ntihistamine Brand or Go	eneric:	
ntihistamine Dose:		



FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN


EPIPEN® (EPINEPHRINE) AUTO-INJECTOR DIRECTIONS

- 1. Remove the EpiPen Auto-Injector from the plastic carrying case.
- 2. Pull off the blue safety release cap.
- 3. Swing and firmly push orange tip against mid-outer thigh.
- 4. Hold for approximately 10 seconds.
- 5. Remove and massage the area for 10 seconds.




AUVI-Q™ (EPINEPHRINE INJECTION, USP) DIRECTIONS

- Remove the outer case of Auvi-Q. This will automatically activate the voice instructions.
- 2. Pull off red safety guard.
- 3. Place black end against mid-outer thigh.
- 4. Press firmly and hold for 5 seconds.
- 5. Remove from thigh.



ADRENACLICK®/ADRENACLICK® GENERIC DIRECTIONS

- 1. Remove the outer case.
- 2. Remove grey caps labeled "1" and "2".
- 3. Place red rounded tip against mid-outer thigh.
- 4. Press down hard until needle penetrates.
- 5. Hold for 10 seconds. Remove from thigh.



OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Note: ONLY Epinephrine and Asthma inhaler will go on a field trip.

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can get worse quickly.

EMERGENCY CONTACTS — CALL 91	OTHER EMERGENCY CONTACTS
RESCUE SQUAD:	NAME/RELATIONSHIP;
DOCTOR:PHONE:	PHONE:
PARENT/GUARDIAN: PHONE:	NAME/RELATIONSHIP:
	PHONE:

PARENT/GUARDIAN AUTHORIZATION SIGNATURE

DATE

PHYSICIAN/HCP AUTHORIZATION SIGNATURE

DATE